

# BRCA1(Ab-1524) Antibody

Catalog No: #21139



Package Size: #21139-1 50ul #21139-2 100ul #21139-4 25ul


## Overview

Product Name	BRCA1(Ab-1524) Antibody
Host Species	Rabbit
Clonality	Polyclonal
Applications	WB IF
Species Reactivity	Hu
Immunogen Type	Peptide-KLH
Target Name	BRCA1
Alternative Names	RNF53


## Application Details

Predicted MW: 220kd
Western blotting: 1:500~1:1000
Immunofluorescence: 1:100~1:200

## Images



Western blot analysis of extracts from 293 and MCF cells using BRCA1(Ab-1524) Antibody #21139.



Immunofluorescence staining of methanol-fixed HeLa cells using BRCA1(Ab-1524) Antibody #21139.

## Descriptions

Immunogen	Peptide sequence around aa.1522~1526 (Y-P-S-Q-E) derived from Human BRCA1.
Specificity	The antibody detects endogenous level of total BRCA1 protein.
Purification	Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific peptide.
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg <sup>2+</sup> and Ca <sup>2+</sup> ), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.
Accession NO.	Swiss-Prot: P38398NCBI Protein: NP_009225.1

## Related Information

The BRCA1-BARD1 heterodimer coordinates a diverse range of cellular pathways such as DNA damage repair, ubiquitination and transcriptional regulation to maintain genomic stability. Acts by mediating ubiquitin E3 ligase activity that is required for its tumor suppressor function. Plays a central role in DNA repair by facilitating cellular response to DNA repair. Required for appropriate cell cycle arrests after ionizing irradiation in both the S-phase and the G2 phase of the cell cycle. Involved in transcriptional regulation of P21 in response to DNA damage. Required for FANCD2 targeting to sites of DNA damage. May function as a transcriptional regulator. Inhibits lipid synthesis by binding to inactive phosphorylated ACACA and preventing its dephosphorylation

Lin SY, et al. (2004) Proc Natl Acad Sci U S A, 101(17): 6484-6489

Zhang J, et al. (2004) Mol Cell Biol; 24(2): 708-718

Beger C, et al. (2001) Proc Natl Acad Sci U S A; 98(1): 130-135

Gardner K, et al. (2001) Breast Cancer Res; 3(1): 11-13

Zheng L, et al. (2001) Proc Natl Acad Sci U S A; 98(17): 9587-9592

**Note:** This product is for in vitro research use only and is not intended for use in humans or animals.