

PAK1/PAK2/PAK3(Ab-423/402/421) Antibody

Catalog No: #21169



Package Size: #21169-1 50ul #21169-2 100ul #21169-4 25ul


Overview

Product Name	PAK1/PAK2/PAK3(Ab-423/402/421) Antibody
Host Species	Rabbit
Clonality	Polyclonal
Applications	WB IHC IF
Species Reactivity	Hu Ms Rt
Immunogen Type	Peptide-KLH
Target Name	PAK1/PAK2/PAK3
Alternative Names	p21-activated kinase 1; PAK-1; p65-PAK; Alpha-PAK;


Application Details

Predicted MW: 61-67 (PAK2) 68-74 (PAK1/3)kd
Western blotting: 1:500~1:1000
Immunohistochemistry: 1:50~1:100
Immunofluorescence: 1:100~1:200


Images



Western blot analysis of extracts from MCF-7 and Hela cells using PAK1/PAK2/PAK3(Ab-423/402/421) Antibody #21169.



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using PAK1/PAK2/PAK3(Ab-423/402/421) Antibody #21169(left) or the same antibody preincubated with blocking peptide(right).



Immunofluorescence staining of methanol-fixed HeLa cells using PAK1/PAK2/PAK3(Ab-423/402/421) Antibody #21169.

## Descriptions

Immunogen	Peptide sequence around aa.421~425/400~404/419~423 (R-S-T-M-V) derived from Human PAK1/PAK2/PAK3.
Specificity	The antibody detects endogenous level of total PAK1/PAK2/PAK3 protein.
Purification	Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific peptide.
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg <sup>2+</sup> and Ca <sup>2+</sup> ), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.
Accession NO.	Swiss-Prot: Q13153/Q13177/O75914NCBI Protein: NP_001122092.1 NP_002568.2 NP_001121638.1

## Related Information

The activated kinase acts on a variety of targets. Likely to be the GTPase effector that links the Rho-related GTPases to the JNK MAP kinase pathway. Activated by CDC42 and RAC1. Involved in dissolution of stress fibers and reorganization of focal complexes. Involved in regulation of microtubule biogenesis through phosphorylation of TBCB. Activity is inhibited in cells undergoing apoptosis, potentially due to binding of CDC2L1 and CDC2L2.

Parsons M, et al. (2005) Mol Cell Biol ; 25:1680-1695.

Jung JH, et al. (2005) J Biol Chem; 280:40025-40031.

Lampugnani MG, et al. (2002) Mol Biol Cell ; 13: 1175-1189

Wang RA, et al. (2002) EMBO J; 21: 5437 - 5447.

**Note:** This product is for in vitro research use only and is not intended for use in humans or animals.