

# 4E-BP1(Ab-45) Antibody

Catalog No: #21216



Package Size: #21216-1 50ul #21216-2 100ul #21216-4 25ul


## Overview

Product Name	4E-BP1(Ab-45) Antibody
Host Species	Rabbit
Clonality	Polyclonal
Applications	WB IHC IF
Species Reactivity	Hu Ms Rt
Immunogen Type	Peptide-KLH
Target Name	4E-BP1
Alternative Names	EIF4EBP1; PHAS-1;


## Application Details

Predicted MW: 18kd
Western blotting: 1:500~1:1000
Immunohistochemistry: 1:50~1:100
Immunofluorescence: 1:100~1:200


## Images



Western blot analysis of extracts from 293 and HepG2 cells using 4E-BP1(Ab-45) Antibody #21216.



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using 4E-BP1(Ab-45) Antibody #21216(left) or the same antibody preincubated with blocking peptide(right).



Immunofluorescence staining of methanol-fixed HeLa cells using 4E-BP1(Ab-45) Antibody #21216.

## Descriptions

Immunogen	Peptide sequence around aa.43~47 (S-T-T-P-G) derived from Human 4E-BP1.
Specificity	The antibody detects endogenous level of total 4E-BP1 protein.
Purification	Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific peptide.
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg <sup>2+</sup> and Ca <sup>2+</sup> ), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.
Accession NO.	Swiss-Prot: Q13541NCBI Protein: NP_004086.1

## Related Information

4E-BP1 encodes one member of a family of translation repressor proteins. The protein directly interacts with eukaryotic translation initiation factor 4E (eIF4E), which is a limiting component of the multisubunit complex that recruits 40S ribosomal subunits to the 5' end of mRNAs. Interaction of this protein with eIF4E inhibits complex assembly and represses translation. This protein is phosphorylated in response to various signals including UV irradiation and insulin signaling, resulting in its dissociation from eIF4E and activation of mRNA translation.

Gingras AC, et al. (1998) *Genes Dev* 12(4): 502-513.

Brugarolas J, et al. (2004) *Genes Dev* 18(23): 2893-2904.

Kumar V, et al. (2000) *EMBO J* 19(5): 1087-1097.

Moody CA, et al. (2005) *J Virol* 79(9): 5499-5506.

Burnett PE, et al. (1998) *Proc Natl Acad Sci U S A* 95(4): 1432-1437.

**Note:** This product is for in vitro research use only and is not intended for use in humans or animals.