

ICAM-1(Phospho-Tyr512) Antibody

Catalog No: #11083



Package Size: #11083-1 50ul #11083-2 100ul #11083-4 25ul



Overview

Product Name	ICAM-1(Phospho-Tyr512) Antibody
Host Species	Rabbit
Clonality	Polyclonal
Applications	WB
Species Reactivity	Hu
Immunogen Type	Peptide-KLH
Target Name	ICAM-1
Modification	Phospho-Tyr512
Alternative Names	ICA1; ICAM1;

Application Details

Predicted MW: 89 92kd
Western blotting: 1:500~1:1000

Images



Descriptions

Immunogen	Peptide sequence around phosphorylation site of tyrosine 512 (K-K-Y(p)-R-L) derived from Human ICAM-1.
Specificity	The antibody detects endogenous level of ICAM-1 only when phosphorylated at tyrosine 512.
Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography using non-phosphopeptide.
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.
Accession NO.	Swiss-Prot: P05362NCBI Protein: NP_000192.2

Related Information

ICAM proteins are ligands for the leukocyte adhesion protein LFA-1 (integrin α -L/ β -2). During leukocyte trans-endothelial migration, ICAM1 engagement promotes the assembly of endothelial apical cups through SGEF and RHOG activation. In case of rhinovirus infection acts as a cellular receptor for the virus.

Greenwood J, et al. (2003) J Immunol; 171(4):2099-2108.

Zhou Z, et al. (2005) Eur J Pharmacol; 513(1-2):1-8.

Chen YH, et al. (2001) J Cell Biochem; 82(3):512-521

Note: This product is for in vitro research use only and is not intended for use in humans or animals.