

Rb(Phospho-Ser795) Antibody

Catalog No: #11130



Package Size: #11130-1 50ul #11130-2 100ul #11130-4 25ul


Overview

Product Name	Rb(Phospho-Ser795) Antibody
Host Species	Rabbit
Clonality	Polyclonal
Applications	WB IHC IF
Species Reactivity	Hu Ms Rt
Immunogen Type	Peptide-KLH
Target Name	Rb
Modification	Phospho-Ser795
Alternative Names	P105-RB; PP105; PP110; RB-1; RB1


Application Details

Predicted MW: 110kd
Western blotting: 1:500~1:1000
Immunohistochemistry: 1:50~1:100
Immunofluorescence: 1:100~1:200


Images



Western blot analysis of extracts from K562 cells untreated(lane 1) or treated with serum(lane 2) using Rb(Phospho-Ser795) Antibody #11130.



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using Rb(Phospho-Ser795) Antibody #11130(left) or the same antibody preincubated with blocking peptide(right).



Immunofluorescence staining of methanol-fixed HeLa cells using Rb(Phospho-Ser795) Antibody #11130.

Descriptions

Immunogen	Peptide sequence around phosphorylation site of serine 795 (P-S-S(p)-P-L) derived from Human Rb.
Specificity	The antibody detects endogenous level of Rb only when phosphorylated at serine 795.
Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography using non-phosphopeptide.
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.
Accession NO.	Swiss-Prot: P06400NCBI Protein: NP_000312.2

Related Information

Key regulator of entry into cell division that acts as a tumor suppressor. Acts as a transcription repressor of E2F1 target genes. The underphosphorylated, active form of RB1 interacts with E2F1 and represses its transcription activity, leading to cell cycle arrest. Directly involved in heterochromatin formation by maintaining overall chromatin structure and, in particular, that of constitutive heterochromatin by stabilizing histone methylation. Recruits and targets histone methyltransferases SUV39H1, SUV420H1 and SUV420H2, leading to epigenetic transcriptional repression. Controls histone H4 'Lys-20' trimethylation. Inhibits the intrinsic kinase activity of TAF1. In case of viral infections, interactions with SV40 large T antigen, HPV E7 protein or adenovirus E1A protein induce the disassembly of RB1-E2F1 complex thereby disrupting RB1's activity.

Fang Sun , Hanjiang Fu , Qin Liu, et al. (2008) Downregulation of CCND1 and CDK6 by miR-34a induces cell cycle arrest. FEBS Letters, 582:1564

Published Papers

Fang Sun, Hanjiang Fu, Qin Liu et al., Downregulation of CCND1 and CDK6 by miR-34a induces cell cycle arrest., FEBS Letters, 582:1564-1568. (2008)

[PMID:18406353](#)

Heidi Braumuller, Thomas Wieder, Ellen Brenner et al., T-helper-1-cell cytokines drive cancer into senescence., NATURE, 494:361-366(2013)

[PMID:23376950](#)

Note: This product is for in vitro research use only and is not intended for use in humans or animals.