Stathmin1(Phospho-Ser25) Antibody

Catalog No: #11224

Package Size: #11224-1 50ul #11224-2 100ul #11224-4 25ul



Overview

Product Name	Stathmin1(Phospho-Ser25) Antibody
Host Species	Rabbit
Clonality	Polyclonal
Applications	WB IHC
Species Reactivity	Hu Ms Rt
Immunogen Type	Peptide-KLH
Target Name	Stathmin1
Modification	Phospho-Ser25
Alternative Names	STMN1; STN1; stathmin


Application Details

Predicted MW: 19kd


Western blotting: 1:500~1:1000

Immunohistochemistry: 1:50~1:100

Images



Western blot analysis of extracts from Jurkat cells untreated or treated with PMA using Stathmin 1(Phospho-Ser25) Antibody #11224.



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using Stathmin 1(Phospho-Ser25) Antibody #11224(left) or the same antibody preincubated with blocking peptide(right).

Descriptions

nunogen	Peptide sequence around phosphorylation site of serine 25 (I-L-S(p)-P-R) derived from Human Stathmin 1.
ecificity	The antibody detects endogenous level of Stathmin 1 only when phosphorylated at serine 25.
ifiction	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates.
	Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho
f	specific antibodies were removed by chromatogramphy using non-phosphopeptide.
mulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02%
1	sodium azide and 50% glycerol.
rage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.
ession NO.	Swiss-Prot: P16949NCBI Protein: NP_001138926.1
mulation :	Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phosphopeptide specific antibodies were removed by chromatogramphy using non-phosphopeptide. Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.0 sodium azide and 50% glycerol. Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.

Related Information

Involved in the regulation of the microtubule (MT) filament system by destabilizing microtubules. Prevents assembly and promotes disassembly of microtubules. Phosphorylation at Ser-16 may be required for axon formation during neurogenesis. Involved in the control of the learned and innate fear Boehm M, et al. (2002) EMBO J 21(13): 3390-3401.

Vadlamudi RK, et al. (2005) Mol Cell Biol 25(9): 3726-3736.

Zilfou JT, et al. (2001) Mol Cell Biol 21(12): 3974-3985.

Biernat J, et al. (2002) Mol Biol Cell 13(11): 4013-4028.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.