

IGF-1R(Phospho-Tyr1280) Antibody

Catalog No: #11302



Package Size: #11302-1 50ul #11302-2 100ul #11302-4 25ul

Overview


Product Name	IGF-1R(Phospho-Tyr1280) Antibody
Host Species	Rabbit
Clonality	Polyclonal
Applications	WB
Species Reactivity	Human
Immunogen Type	Peptide-KLH
Target Name	IGF-1R
Modification	Phospho-Tyr1280
Alternative Names	IGFR; CD221; IGFIR

Application Details

Predicted MW: 200 95kd

Western blotting: 1:500~1:1000

Images



Western blot analysis of extract from 293 cells using IGF-1R(Phospho-Tyr1280) Antibody #11302.

Descriptions

Immunogen	Peptide sequence around phosphorylation site of tyrosine 1280 (S-F-Y(p)-Y-S) derived from Human IGF-1R
Specificity	The antibody detects endogenous levels of IGF-1R only when phosphorylated at tyrosine 1280.
Purification	Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific peptide.
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.
Accession NO.	Swiss-Prot: P08069NCBI Protein: NP_000866.1

Related Information

This receptor binds insulin-like growth factor 1 (IGF1) with a high affinity and IGF2 with a lower affinity. It has a tyrosine-protein kinase activity, which is necessary for the activation of the IGF1-stimulated downstream signaling cascade. When present in a hybrid receptor with INSR, binds IGF1. Ref.19 shows that hybrid receptors composed of IGF1R and INSR isoform Long are activated with a high affinity by IGF1, with low affinity by IGF2 and not significantly activated by insulin, and that hybrid receptors composed of IGF1R and INSR isoform Short are activated by IGF1, IGF2 and insulin. In contrast, Ref.21 shows that hybrid receptors composed of IGF1R and INSR isoform Long and hybrid receptors composed of IGF1R and INSR isoform Short have similar binding characteristics, both bind IGF1 and have a low affinity for insulin.

Adams, T.E. et al. (2000) Cell. Mol. Life Sci. 57, 1050-1093

Baserga, R. et al. (2000) Oncogene 19, 5574-5581

Scheidegger, K.J. et al. (2000) J. Biol. Chem. 275, 38921-38928.

O'Connor R, et al. Mol Cell Biol 1997 Jan; 17(1): 427-35

Note: This product is for in vitro research use only and is not intended for use in humans or animals.