Product Datasheet

Shc1(Phospho-Tyr427) Antibody

Catalog No: #11317



Package Size: #11317-1 50ul #11317-2 100ul #11317-4 25ul

Overview

Product Name	Shc1(Phospho-Tyr427) Antibody
Host Species	Rabbit
Clonality	Polyclonal
Applications	WB IHC IF
Species Reactivity	Hu Ms Rt
Immunogen Type	Peptide-KLH
Target Name	Shc1
Modification	Phospho-Tyr427
Alternative Names	SH2 domain protein C1; SHC; SHC-transforming protein 1; SHCA; Src homology 2
	domain-containing-transforming protein C1

Application Details

Predicted MW: 46 52 kd	
Western blotting: 1:500~1:1000	
Immunohistochemistry: 1:50~1:100	
Immunofluorescence: 1:100~1:200	

Images



Western blot analysis of extracts from 293 cells untreated or treated with PMA using Shc1(Phospho-Tyr427) Antibody #11317.



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using Shc1(Phospho-Tyr427) Antibody #11317(left) or the same antibody preincubated with blocking peptide(right).



Descriptions	
Immunogen	Peptide sequence around phosphorylation site of tyrosine 427 (P-S-Y(p)-V-N derived from Human Shc1.
Specificity	The antibody detects endogenous level of Shc1 only when phosphorylated at tyrosine 427.
Purifiction	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatogramphy using non-phosphopeptide.
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.
Accession NO.	Swiss-Prot: P29353NCBI Protein: NP_001123512.1

Related Information

Signaling adapter that couples activated growth factor receptors to signaling pathway. Isoform p46Shc and isoform p52Shc, once phosphorylated, couple activated receptor tyrosine kinases to Ras via the recruitment of the GRB2/SOS complex and are implicated in the cytoplasmic propagation of mitogenic signals. Isoform p46Shc and isoform p52Shc may thus function as initiators of the Ras signaling cascade in various non-neuronal systems. Isoform p66Shc does not mediate Ras activation, but is involved in signal transduction pathways that regulate the cellular response to oxidative stress and life span. Isoform p66Shc acts as a downstream target of the tumor suppressor p53 and is indispensable for the ability of stress-activated p53 to induce elevation of intracellular oxidants, cytochrome c release and apoptosis. The expression of isoform p66Shc has been correlated with life span Trampont P, et al. (2006) Mol Cell Biol; 26(23): 9035-9044.

Patrussi L, et al. (2005) Oncogene; 24(13): 2218-2228

van der Geer P, et al. (1996) Curr Biol ; 6(11): 1435-1444

Note: This product is for in vitro research use only and is not intended for use in humans or animals.